

RESEARCH ARTICLE

Is the removal of aboveground shrub biomass an effective technique to restore a shrub-encroached grassland?

Yanshu Liu¹, Zhongjie Shi¹, Liyan Gong¹, Richun Cong¹, Xiaohui Yang^{1,2}, David J. Eldridge³ 

Encroachment of woody plants into grasslands is a global phenomenon that has substantial impacts on pastoral productivity and ecosystem services. Over the past half century, pastoralists and land management agencies have explored various options to control woody plants in order to improve ecosystem services in shrub-encroached grasslands. We examined the effectiveness of controlling the encroachment of the shrub *Caragana microphylla* into grassland in Inner Mongolia, China. We cut and removed all of the aboveground biomass from 450 shrubs, predicting that the effectiveness of this technique to control shrubs would depend on shrub morphology. Specifically, we expected that larger shrubs with more biomass would be more difficult to kill by cutting than smaller shrubs. A year after treatment, we found that cutting killed only 11% of the 450 treated shrubs, and of these, three-quarters of the locations that they occupied reverted to grasses and one-quarter to bare soil. Shrubs that survived the cutting treatment produced more stems and leaf biomass, and therefore had a greater leaf to stem ratio. Shrubs that died after cutting had a lower crown area and basal area, and less stem biomass than shrubs that resprouted within 12 months of cutting. There were no effects of shrub height on the fate of treated shrubs. Cutting had no effect on understory plant cover or richness, but reproductive plants were taller under shrubs that were not cut. Overall, our study showed that removing aboveground shrub biomass by cutting is an ineffective technique for “restoring” the original grassland community unless shrubs are very small. Strategic targeting of small shrubs would be a more effective technique for controlling the spread of *C. microphylla* in the long term.

Key words: *Caragana microphylla*, encroachment, grassland condition, mechanical removal

Implications for Practice

- Removal of the aboveground biomass of *Caragana microphylla* shrubs by cutting is a relatively ineffective technique for managing shrub densities if the aim is to “restore” the original grassland community.
- The value of cutting depends on whether soil hummocks supporting shrubs provide greater short-term production for grazing.
- Land managers wishing to control *C. microphylla* by cutting (e.g. mowing) should concentrate on sites with small shrubs where treatment is more effective.

Introduction

Woody encroachment is a global phenomenon characterized by an increase in the cover and/or density of woody plants at the expense of herbaceous species, often grasses (Van Auken 2000). The causal mechanisms of encroachment are multifaceted, and relate to changes in fire regimes, overgrazing by domestic livestock, land use disturbance, and increasing concentrations of atmospheric CO₂ (Van Auken 2000; Eldridge et al. 2011; Archer & Predick 2014). At a global scale, woody encroachment is thought to affect a large area of the world’s drylands, a biome that supports about 40% of the global human population, many

of whom rely on pastoral production for their livelihoods (Pravali 2016). Over the past half century, practitioners have invested substantial funds to manage the effects of woody encroachment in the Americas, Australia, and Africa (Paynter & Flanagan 2004; Noble & Walker 2006; Isaacs et al. 2013; Archer & Predick 2014). For example, the Restore New Mexico Program aims to reverse economic loss caused by woody encroachment (Zavaleta et al. 2001; Hamilton et al. 2004). Control of woody plants is often motivated by the need to increase pastoral production and therefore farmer livelihoods, though other reasons can be to alter habitat for threatened plants and animals or to reduce the risk of wildfire (Huggett et al. 2008). Removal is based on the often perverse assumption that shrubs and herbaceous plants

Author contributions: YL, ZS conceived and designed the research; YL, LG, RC performed the experiments; YL, DJE analyzed the data; YL, DJE, XY wrote and edited the manuscript.

¹Institute of Desertification Studies, Chinese Academy of Forestry, Beijing, 100091, China

²Address correspondence to Xiaohui Yang, email yangxh@caf.ac.cn

³Centre for Ecosystem Science, School of Biological, Earth and Environmental Sciences, University of New South Wales, Sydney, New South Wales 2052, Australia

are incompatible, and that removal of woody plants is necessary to increase pastoral production (Eldridge et al. 2013).

The physical removal of woody plants is a popular and widely used technique for reducing woody plant density and cover (Archer et al. 2011). A wide range of physical techniques have been used, with variable success. For example, blade (root) plowing is used in Australia to reduce extensive stands of native shrub species such as *Eremophila* and *Dodonaea* (Robson 1995). In the Americas, physical methods such as roller-chopping have been trialed to reduce the density and cover of woody plants (Adema et al. 2004). Although different woody plant management techniques can be applied singularly or with other techniques, results to date suggest that one-off treatments are generally short lived (Archer et al. 2011). Results might also depend on the type of treatment. For example, mechanical treatments that covered the severed roots of the shrub *Eremophila sturtii* have been shown to be more effective at killing shrubs than those that leave the roots exposed (Wiedemann & Kelly 2001). In China, some pastoralists have experimented with mowing shrubs in the expectation that this will reduce shrub growth rate and stimulate the growth of perennial grasses. However, to our knowledge, we know of no studies that have specifically tested the efficacy of such cutting treatments and whether the effectiveness of cutting varies with differences in shrub attributes such as size. Results might also depend on the type of treatment. For example, mechanical treatments that cover the severed roots of the shrub *Eremophila sturtii* have been shown to be more effective at removal than those that leave the roots exposed (Wiedemann & Kelly 2001).

The response of woody plants to different treatments depends on the identity of the species and environmental conditions. The success of any woody plant removal program, therefore, will depend on the mechanisms by which plants reduce the effects of herbivory to maximize survival and growth under different disturbance and environmental regimes (Meloche & Diggle 2001). These include diverting resources from the production of reproductive structures, which are carbohydrate expensive, to defense structures such as thorns (Zhang et al. 2006) or increases in anti-herbivore leaf phenols (Simms & Rausher 1987), resprouting from epicormic buds, or reestablishing from the soil seed bank (Götmark et al. 2016). Plants that invest in the production of multiple stems may be shorter (Midgley 1996; Paula & Pausas 2011), but removal of aboveground material could stimulate resprouting species to produce more shoots per individual than occurred prior to disturbance (Hermann et al. 2012). This has been observed in the conspecific *Caragana korshinskii* where total removal of aboveground material stimulated long side shoots, allowing shrubs to maximize vertical structure as quickly as possible in order to escape further herbivory (Fang et al. 2006a). Finally, larger shrubs may be able to allocate more resources to roots to sustain high growth rates (Paula & Pausas 2011). Thus, larger shrubs are likely to have a greater root to shoot ratio, allowing them to resprout more easily after defoliation or physical disturbance because they can more readily allocate resources to aboveground compartments.

Caragana microphylla Lam. is a perennial xerophytic shrub from the family Fabaceae, and currently occupies more than

5 million ha of China's northern grassland (Zhang et al. 2006; Peng et al. 2013a). Although *microphylla* is a natural component of these grasslands, the cover and abundance of this plant has increased markedly over the past half century. Considerable research suggests that encroachment of *Caragana* results from overgrazing by domestic livestock, which reduces competition for soil moisture from perennial grasses (Hester et al. 1996; Zhang et al. 2018), and increases in atmospheric CO₂ concentrations (Xiong et al. 2003; Christensen et al. 2004). Once established, *Caragana* is relatively resistant to grazing (Xiong et al. 2003), leading to its widespread persistence. Encroachment is enhanced by the fact that *Caragana* is highly temperature- and drought-tolerant, and therefore has the capacity to rapidly increase in density under adverse conditions (Cao et al. 2000; Chen et al. 2015). Although encroachment of *C. microphylla* into grasslands could provide shelter for livestock and increase landscape-level biodiversity by facilitating understory plants (Liu 1991), the grazing value of the previously grass-dominant steppe for livestock has declined (Xiong et al. 2003; Xiong & Han 2005; Peng et al. 2013b). The presence of *Caragana* also increases the difficulty of hay production by mowing (Liu 1991). Young *C. microphylla* plants can be controlled by fire, which will also reduce bud formation on mature plants (Li & Jiang 1994; Lin & Bai 2010). However, current government policy prevents the widespread use of fire or mechanical methods to control to remove shrubs.

In this study, we examined the effectiveness of controlling the encroachment of *C. microphylla* by removing the aboveground compartment of the shrubs by cutting. Specifically, we asked whether shrub density can be reduced by removing the aboveground biomass and thus whether cutting is a potential technique for managing the encroachment of *C. microphylla* into *Leymus chinensis* (Trin.) Tzvel and *Stipa grandis* P. Smirn grassland. A potential state-and-transition model for *C. microphylla* suggests a transition from State A (shrubland) to State B (bare soil) with removal of aboveground shrub biomass (Fig. 1) and a further transition to State C (grassland) dominated by grasses and/or forbs triggered by removal of the competitive effects of the shrubs. Equally likely, however, is a transition to State D (recovering shrubland) depending on the extent of shrub recovery. The question of what determines shrub survival after biomass removal by cutting is poorly known. This lack of information limits our capacity to recommend cost-effective management options to control shrubs in grasslands or to reduce their density. We expected that the effectiveness of cutting would vary with shrub morphology (Fang et al. 2006b). For example, taller plants and those with a greater basal area and more biomass would be more likely to resprout after cutting because they can allocate resources to recover from disturbance (Massi & Franco 2016) or overcompensate with greater bud growth, such as occurs with bitterbrush (*Purshia tridentata*) a dryland shrub (Bilbrough & Richards 1993). In contrast, smaller shrubs with smaller canopies and less biomass might be expected to be more susceptible to cutting because resources would be rapidly depleted by cutting, and shrubs would be less likely to resprout (Fig. 1).




Figure 1. Conceptual model of the potential states and transitions within a *Caragana microphylla* encroached grassland–shrubland matrix. The state-and-transition model captures potential scenarios that might arise after all aboveground material is removed from the encroaching shrubs by cutting. In our model, numbers indicate transitions and letters indicate states. The starting position in our model is an encroached grassland (State A) with scattered but diminishing grasses. Both the understory shrub environment and the interspaces likely vary in plant composition depending on shrub size and abiotic conditions, and, as suggested in our study, in the heights of plants with reproductive structures. Transition 1 is achieved by shrub biomass removal, leading to a predicted temporary state (State B), which, in our study, shifted rapidly to State D within 12 months due to the resprouting ability of *Caragana*. Few plants shifted to State C (Transition 2), which is likely driven by rainfall and a relaxation of grazing intensity. Based on previous studies, we would expect State C to move to State D (Transition 5) with overgrazing and increased atmospheric CO₂ (Xiong et al. 2003; Zhang et al. 2006), but the time frame is likely variable. Transition 4 is likely to occur over less than a decade in the absence of other interventions.

Specifically, we tested whether the extent to which cut shrubs (State B; Fig. 1) transition (via Transition 2) to State C (grasses present, shrubs absent) or State D (via Transition 3, shrubs resprouting with sparse grasses), and posit that larger shrubs would resprout (Transition 3) whereas smaller shrubs would succumb to cutting (Transition 2). We tested our prediction by comparing differences in size-related plant traits between individuals that either survived or died following the cutting treatment. We then compared the pretreatment, size-related traits of plants that either died or survived (resprouted) to test our hypothesis that larger shrubs were more resistant to the removal of their aboveground biomass.

Methods

Study Site

This study was carried out in the typical steppe of Inner Mongolia, China (44.372°N, 44.147°E), a central region of the Eurasian steppe (Bai et al. 2008). Winters are cold and dry (average daily temperature -22°C) and summers mild and humid (average 19°C) with an average annual rainfall of about 290 mm, 60–80% falling in the growing season (June–September). The soil is predominantly a calcareous chestnut soil (Calcic Chernozem) according to the IUSS Working Group WRB (2006) and the slopes are $<1\%$. The plant community is dominated by the perennial tussock grasses *Stipa grandis* and the perennial rhizomatous grass *Leymus chinensis* (Bai et al. 2004). A more detailed description of the plant community and response of dominant species to grazing at this site is given in Liu et al. (2011). The area also shows evidence of encroachment by the thorny leguminous shrub *Caragana microphylla* due to half a century of overgrazing and intense landuse change. *C. microphylla* now occurs in

many places within a matrix of grass and shrub patches (Peng et al. 2013a).

In 2017, we established the experiment in a 67 ha grassland site with a uniform distribution of *C. microphylla* shrubs. The site was moderately grazed prior to livestock exclusion fencing. In July 2017, we randomly selected fifty 100-m² plots (each 10 m \times 10 m) within this site and marked their location. Within each plot we marked the position of every *C. microphylla* shrub and its hummock (the slightly elevated area around each plant) and, for each shrub, measured the height, crown diameter, and basal diameter in two directions, and counted the number of stems before cutting off all of the aboveground material from each shrub. Shrub density averaged 906 ± 55 shrubs/ha (mean \pm SE) and cover was $5.9 \pm 0.6\%$. In July 2018, 12 months after shrub biomass removal, we relocated the plots and individual shrubs, and made the same measurements as prior to biomass removal. Shrubs were recorded as resprouted or dead on bare soil or dead with grasses. Mean shrub density in July 2018 was 802 ± 53 shrubs/ha and cover $4.7 \pm 0.5\%$. For each shrub, we separated leaf material, new stems, and old stems, and dried the material until constant weight, then weighed the material.

To assess understory vegetation, we randomly selected 20 of the plots where shrubs had been cut, and an additional 20 control plots where shrubs had not been cut. Within each of these plots we measured total plant cover, and the height of up to five vegetative (without reproductive structures) and reproductive plants within two microsites: under shrub canopies and in the open.

Statistical Analyses

We tested for differences in shrub attributes and understory plant attributes (e.g. plant cover, richness) before and after cutting

using linear models. For shrub attributes, our model had 50 blocks (plots) and two treatments (before and after biomass removal), and we tested the effect of biomass removal using the block by treatment residual term. Analyses were similar for understory plant attributes except that the model was a split-plot with two strata, the first examined Treatment (cutting vs. control), and the second Microsite (shrub vs. open) and its interaction with Treatment. Differences in mean values were compared using least significant difference testing. Analyses were conducted after testing for normality and homogeneity of variance (Levene's test) using the Minitab statistical software. The leaf to stem ratio was calculated based on biomass.

Results

We recorded a total of 450 shrubs in our study site across an area of 5 ha (90 plants/ha). Of the 450 shrubs, 89% (401 shrubs) regrew. Most (84%, 377) of the 401 shrubs regrew completely, and the remaining 24 exhibited partial regrowth. Hummocks with dead shrubs either reverted to bare soil (12) or were colonized by grasses (37; Fig. 2). We also recorded an additional 16 new shrub recruits across the 5-ha sites measured.

Effects of Shrub Biomass Removal Treatment on Shrub Attributes

Twelve months after biomass removal, surviving shrubs, that is those showing partial or complete regrowth, had markedly different morphologies compared with pretreatment. For example, surviving shrubs were 16% shorter ($F_{[1,399]} = 31.9, p < 0.001$), had 67% smaller crown areas ($F_{[1,399]} = 31.9, p < 0.001$), 64% smaller basal area ($F_{[1,399]} = 31.9, p < 0.001$), less total biomass ($F_{[1,399]} = 31.9, p < 0.001$), and stem biomass ($F_{[1,399]} = 31.9, p < 0.001$; Fig. 3). Our shrub treatment also stimulated the production of stems by 124% ($F_{[1,399]} = 31.9, p < 0.001$), total shrub leaf biomass by 84% ($F_{[1,399]} = 31.9, p < 0.001$), and, consequently, increased the leaf:stem ratio by 178% ($F_{[1,399]} = 31.9, p < 0.001$; Fig. 3).

Do Shrub Attributes Explain Their Susceptibility to Biomass Removal?

Shrubs growing on hummocks that had either reverted to bare soil or grass tended to have fewer stems per plant ($F_{[3,446]} = 14.59, p < 0.001$). Shrubs growing on hummocks reverting to bare soil tended to have lower crown ($F_{[3,446]} = 11.11, p < 0.001$) and basal areas ($F_{[3,446]} = 12.02, p < 0.001$) while those that reverted to grasses tended to have significantly less stem ($F_{[3,446]} = 6.10, p < 0.001$) and total biomass ($F_{[3,446]} = 6.70, p < 0.001$) than those that either resprouted totally or partially following treatment (Fig. 4). There were no apparent effects of shrub height ($p = 0.13$) or leaf:stem ratio ($p = 0.29$) on the fate of shrubs to the cutting treatment (Fig. 4).




Figure 2. The number of *Caragana microphylla* shrubs recorded 12 months after the shrub cutting treatment.

Effects of Cutting on Understory Plants

We found no effect of cutting on either plant cover or plant richness ($p > 0.08$), but plant cover ($F_{[1,18]} = 24.2, p = 0.001$) and richness ($F_{[1,18]} = 18.2, p < 0.001$) were significantly lower, overall, beneath shrubs than in the interspaces (Fig. 5). The heights of vegetative plants did not vary with treatment or microsite ($p > 0.33$). The height of reproductive plants, however, was significantly greater under uncut shrubs than under cut shrubs, irrespective of microsite (treatment by microsite interaction: $F_{[1,18]} = 6.89, p = 0.017$; Fig. 5). Plant attributes were consistently twice as variable under shrubs than in the open across both treatments. Plant richness was more variable under shrubs, but only at cut sites (Table 1). There was little variability in the heights of vegetative and reproductive plants in relation to cutting treatment or microsite.

Discussion

In our study, cutting killed only 11% of treated shrubs, and only 6% of hummocks where shrubs were treated reverted to grasses. A one-off removal of the aboveground biomass of *Caragana microphylla* shrubs was not an effective method to meet our management objective of controlling shrubs and promoting grass recovery. The tendency of *C. microphylla* to recover after the removal of all aboveground biomass (e.g. through grazing) is likely related to its ability to resprout, and perhaps fix nitrogen. The ability to produce nitrogen may increase the tolerance of *C. microphylla* to herbivory as it resprouts, as has been shown in N-producing forbs and grasses (Watson & Ward 1970). *C. microphylla* is known to resprout from subsurface shoots and axial tillers in response to shoot removal (Fang et al. 2006b).

Removal of stems and leaves of *Caragana* will likely reduce growth rates, given the strong links between measures of leaf gas exchange rate (such as net photosynthetic rate, stomatal conductance, and transpiration rate) and the relative growth rate of *Caragana* spp. (Ma et al. 2016). Further, a greater root to shoot ratio should allow plants to absorb water and




Figure 3. Mean (+SE) of different attributes of *Caragana microphylla* shrubs before and after the removal of aboveground shrub biomass. Different superscripts indicate a significant difference in the mean values at $p < 0.05$.

nutrients more efficiently, but this has been shown not to be the case in *C. microphylla*. Vegetation removal by cutting or herbivory may allow shrubs to invest more resources in defenses against stem removal, thereby retarding the growth of individual plants, particularly young plants (Zhang et al. 2006). An alternative mechanism to account for resprouting following biomass removal could be related to reduced levels of biological control from herbivorous insects that are removed when the plant is defoliated.

Many methods of shrub control such as root plowing or roller-chopping involve substantial soil disturbance, and this physical disturbance typically initiates resprouting by activating the growth of epicormic buds (Wiedemann & Kelly 2001; Fulbright et al. 2013). In our study, however, removal of the aboveground shrub biomass involved little or no soil disturbance. Cutting is more likely to maintain the integrity the soil surface, with fewer effects on surface litter and therefore soil seed banks than treatments that disturb the surface (Eldridge & Robson 1997). Notwithstanding the lower levels of disturbance compared with root plowing (e.g. Daryanto & Eldridge 2010; Smith et al. 2013), cutting was not an effective method for

either removing shrubs or improving understory plant cover and richness.

Our results indicated that the effectiveness of shrub cutting was greater for smaller plants with shorter stems and less biomass. Smaller plants are likely younger, suggesting two things. First, control is more effective on younger, smaller plants that will be more difficult to detect if they are below the size of grasses. Pastoralists therefore need to be vigilant to detect small plants before they reach a size that is unresponsive to physical removal. Second, larger shrubs would need to be controlled by alternative methods, which might include herbicide, or may require follow-up treatment of smaller resprouting individuals. However, we found no evidence to suggest any effect of plant height on the fate of treated plants. This could be due to the fact that there were no clear allometric relationships between height and other size measurements, suggesting that height might not be a good indicator of shrub age, given that the shoots can assume a pronounced prostrate habit.

Shrubs support a range of important ecosystem services such as habitat provision, carbon sequestration, and nutrient cycling (Eldridge et al. 2011; Daryanto et al. 2019). A recent global




Figure 4. Mean (+SE) values of attributes of *Caragana microphylla* shrubs in relation to their fate after cutting. Different superscripts indicate a significant difference in the mean values at $p < 0.05$. There were no differences in leaf:stem ratio or shrub height among the four fate classes.

meta-analyses of shrub removal showed that increases in forage production were relatively short-lived (Daryanto et al. 2019). Persistence of *C. microphylla* after cutting, therefore, does not necessarily mean that this treatment is entirely ineffective for either pastoral production or biodiversity conservation given potential short-term benefits of either increases in production or plant diversity. The extent to which *C. microphylla* competes with or facilitates understory species is poorly known (Zhang et al. 2018). Although shrubs are known to compete with understory protégé species for resources, they also enhance species richness via positive (facilitatory) interactions by ameliorating environmental conditions and protecting protégé species from grazing (Smit et al. 2009; Soliveres et al. 2016). The ability of shrubs to facilitate understory protégé species might increase

with shrub age, and therefore size (Zhao et al. 2007). Thus, shrubs resprouting after complete removal of all aboveground material will likely be less effective at facilitating protégé species; many of which are likely palatable to livestock. Physical removal therefore may produce unintended consequences by reducing the capacity of shrubs to facilitate pastorally important understory species.

Our study is based on an assessment of understory locations of individual shrubs and their paired interspaces. Despite not sampling at broader spatial scales (e.g. patches of shrubs, communities), the state-and-transition model identifies the likely scenarios that might emerge at these larger spatial scales. Scaled-up results suggest that within 12 months, State B moves to State D (Transition 3), which is characterized by a substantial




Figure 5. Mean (+SE) values for selected attributes of understory plants in the open and beneath the canopy of *Caragana microphylla* shrubs for both uncut (control) and cut treatments. Different superscripts indicate significant differences at $p < 0.05$.

Table 1. Mean variability (coefficient of variation %) in understory plant cover (%), plant richness (number of species), and heights (cm) of vegetative and reproductive plants in the open and beneath shrubs for sites ($n = 10$) where shrubs were either treated (cut) or untreated (uncut).

Attribute	Uncut		Cut	
	Open	Shrub	Open	Shrub
Plant cover (%)	16.4	31.2	18.9	36.9
Plant richness (number of species)	20.4	24.0	23.1	44.9
Height of vegetative plants (cm)	23.1	28.4	21.2	19.1
Height of reproductive plants (cm)	13.5	15.8	11.5	22.1

number of resprouting shrubs, with few shrubs (8%) reverting to the preferred State C (Fig. 1). Our results also suggest that Transition 2 (State A to State C) is highly dependent upon shrub traits, and likely restricted to shrubs with smaller crowns and basal area, less biomass, and fewer shoots per shrub. Given the short duration of our study, we can only speculate upon the potential transition to State D (Transition 5). State D is likely to be relatively transient, with movement from State C to A (via D) occurring relatively rapidly, and exacerbated by overgrazing, which removes competition from perennial grasses (Hester et al. 1996; Zhang et al. 2018). Our results also suggest that progression to State D has no significant effects on understory plant cover or richness, at least in the short term. However, we found some evidence that reproductive plants might be affected by shrub cutting, if a decline in height is indicative of fewer inflorescences and therefore a reduced potential to produce viable seed (McGrath et al. 2010). Our data also indicate

lower heterogeneity in plant attributes with shrub cutting, which could lead to long-term declines in ecosystem functions. Overall, therefore, any decision to remove shrubs will be highly context-dependent, varying with societal goals for shrublands and/or the need to maintain a healthy, productive and diverse plant community. Land managers, therefore, need to be cognizant of the many values of *C. microphylla* before undertaking control methods designed to increase grass production.

Acknowledgments

This study was supported by the Fundamental Research Funds for the Central Non-profit Research Institution of CAF (YBB2016ZD010), the National Key R&D Program of China (2016YFC0500804), and the National Science Foundation of China (31200350; 41471029; 41271033; 31600581). We thank Professor Yongfei Bai and Associate Professor Dima Chen for their help with experimental design. We thank the staff at the Inner Mongolia Grassland Ecosystem Research Station for their help with field work.

LITERATURE CITED

- Adema EO, Buschiazzo DE, Babinec FJ, Rucci TE, Hermida VFG (2004) Mechanical control of shrubs in a semiarid region of Argentina and its effect on soil water content and grassland productivity. *Agricultural Water Management* 68:185–194
- Archer SR, Predick KI (2014) An ecosystem services perspective on brush management: research priorities for competing land-use objectives. *Journal of Ecology* 102:1394–1407

- Archer S, Davies KW, Fulbright TE, McDaniel KC, Wilcox BP, Predick K, Briske DD (2011) Brush management as a rangeland conservation strategy: a critical evaluation. Pages 105–170. In: Briske DD (ed) Conservation benefits of rangeland practices. US Department of Agriculture Natural Resources Conservation Service, Washington D.C.
- Bai YF, Han XG, Wu JG, Chen ZZ, Li LH (2004) Ecosystem stability and compensatory effects in the Inner Mongolia grassland. *Nature* 431:181–184
- Bai Y, Wu J, Xing Q, Pan Q, Huang J, Yang D, Han X (2008) Primary production and rain use efficiency across a precipitation gradient on the Mongolia plateau. *Ecology* 89:2140–2153
- Bilbrough CJ, Richards JH (1993) Growth of sagebrush and bitterbrush following simulated winter browsing—mechanisms of tolerance. *Ecology* 74:481–492
- Cao C, Jiang D, Alamsa Luo Y, Kou Z (2000) Ecological process of vegetation restoration in *Caragana microphylla* sand-fixing area. *Chinese Journal of Applied Ecology* 11:349–354 (in Chinese)
- Chen L, Li H, Zhang P, Zhao X, Zhou L, Liu T, Hu H, Bai Y, Shen H, Fang J (2015) Climate and native grassland vegetation as drivers of the community structures of shrub-encroached grasslands in Inner Mongolia, China. *Landscape Ecology* 30:1627–1641
- Christensen L, Coughenour MB, Ellis JE, Chen ZZ (2004) Vulnerability of the Asian typical steppe to grazing and climate change. *Climatic Change* 63:351–368
- Daryanto S, Eldridge DJ (2010) Plant and soil surface responses to a combination of shrub removal and grazing in a shrub-encroached woodland. *Journal of Environmental Management* 91:2639–2648
- Daryanto S, Fu B, Zhao W (2019) Evaluating the use of fire to control shrub encroachment in global drylands: a synthesis based on ecosystem service perspective. *Science of the Total Environment* 648:285–292
- Eldridge DJ, Robson AD (1997) Bladeploughing and enclosure influence soil properties in a semi-arid Australian woodland. *Journal of Range Management* 50:191–198
- Eldridge DJ, Bowker MA, Maestre FT, Roger E, Reynolds JF, Whitford WG (2011) Impacts of shrub encroachment on ecosystem structure and functioning: towards a global synthesis. *Ecology Letters* 14:709–722
- Eldridge DJ, Soliveres S, Bowker MA, Val J (2013) Grazing dampens the positive effects of shrub encroachment on ecosystem functions in a semi-arid woodland. *Journal of Applied Ecology* 50:1028–1038
- Fang X, Wang X, Li H, Chen K, Wang G (2006a) Responses of *Caragana korshinskii* to different aboveground shoot removal: combining defence and tolerance strategies. *Annals of Botany* 98:203–211
- Fang X, Yuan J, Wang G, Zhao Z (2006b) Fruit production of shrub, *Caragana korshinskii*, following above-ground partial shoot removal: mechanisms underlying compensation. *Plant Ecology* 187:213–225
- Fulbright TE, Alejandro Lozano-Cavazos E, Ruthven DC III, Lite AR (2013) Plant and small vertebrate composition and diversity 36–39 years after root plowing. *Rangeland Ecology & Management* 66:19–25
- Götmark F, Götmark E, Jensen AM (2016) Why be a shrub? A basic model and hypotheses for the adaptive values of a common growth form. *Frontiers in Plant Science* 7:1095
- Hamilton WT, McGinty A, Ueckert DN, Hanselka CW, Lee MR (2004) Brush management: past, present, future. Texas A&M University Press, College Station, Texas
- Hermann J-M, Haug S, Pillar VD, Pfadenhauer J (2012) Shrubs versus “gullivers”: woody species coping with disturbance in grasslands. *Plant Ecology* 213:1757–1768
- Hester AJ, Mitchell FJG, Kirby KJ (1996) Effects of season and intensity of sheep grazing on tree regeneration in a British upland woodland. *Forest Ecology and Management* 88:99–106
- Huggett RJ Jr, Abt KL, Shepperd W (2008) Efficacy of mechanical fuel treatments for reducing wildfire hazard. *Forest Policy and Economics* 10:408–414
- Isaacs L, Somers MJ, Dalerum F (2013) Effects of prescribed burning and mechanical bush clearing on ungulate space use in an African savannah. *Restoration Ecology* 21:260–266
- IUSS Working Group WRB (2006) World reference base for soil resources 2006. 2nd edition. World Soil Resources Reports No. 103. FAO, Rome, Italy
- Li Z, Jiang Q (1994) The effects of the fire on the nutrient states of steppe soil. *Acta Scientiarum Naturalium Universitatis Neimenggu* 25:444–449
- Lin Y, Bai Y (2010) Responses of aboveground net primary production and population structure of *Caragana microphylla* to prescribed burning in a typical steppe of Inner Mongolia. *Acta Prataculturae Sinica* 19:170–178
- Liu SR (1991) Effects of *Caragana microphylla* Lam. shrub on steppe vegetation in Xilinguole. *Inner Mongolia Prataculture* 3:30–34 (in Chinese)
- Liu Y, Pan Q, Liu H, Bai Y, Simmons M, Dittert K, Han X (2011) Plant responses following grazing removal at different stocking rates in an Inner Mongolia grassland ecosystem. *Plant and Soil* 340:199–213
- Ma F, Na X, Xu T (2016) Drought responses of three closely related *Caragana* species: implication for their vicarious distribution. *Ecology and Evolution* 6:2763–2773
- Massi KG, Franco AC (2016) How does resprouting response differ among three species of savanna trees and in relation to plant size? *Acta Botanica Brasilica* 30:693–699
- McGrath S, Hodkinson TR, Charles TM, Zen DG, Barth S (2010) Variation in inflorescence characters and inflorescence development in ecotypes and cultivars of *Lolium perenne* L. *Grass and Forage Research* 65:398–409
- Meloche CG, Diggle PK (2001) Proliferation, architectural complexity, and developmental flexibility in *Acomastylis rossii* (Rosaceae). *American Journal of Botany* 88:980–991
- Midgley JJ (1996) Why the world's vegetation is not totally dominated by resprouting plants; because resprouters are shorter than reseeders. *Ecography* 19:92–95
- Noble JC, Walker P (2006) Integrated shrub management in semi-arid woodlands of eastern Australia: a systems-based decision support model. *Agricultural Systems* 88:332–359
- Paula S, Pausas JG (2011) Root traits explain different foraging strategies between resprouting life histories. *Oecologia* 165:321–331
- Paynter Q, Flanagan GJ (2004) Integrating herbicide and mechanical control treatments with fire and biological control to manage an invasive wetland shrub, *Mimosa pigra*. *Journal of Applied Ecology* 41:615–629
- Peng H-Y, Li X-Y, Li G-Y, Zhang Z-H, Zhang S-Y, Li L, Zhao G-Q, Jiang Z-Y, Ma Y-J (2013a) Shrub encroachment with increasing anthropogenic disturbance in the semiarid Inner Mongolian grasslands of China. *Catena* 109:39–48
- Peng H, Li X, Tong S (2013b) Effects of shrub encroachment on biomass and biodiversity in the typical steppe of Inner Mongolia. *Acta Ecologica Sinica* 33:7221–7229
- Pravalié R (2016) Drylands extent and environmental issues. A global approach. *Earth-Science Reviews* 161:259–278
- Robson AD (1995) The effects of grazing exclusion and blade-ploughing on semi-arid woodland vegetation in North-Western New South Wales over 30 months. *The Rangeland Journal* 17:111–117
- Simms EL, Rausher MD (1987) Costs and benefits of plant resistance to herbivory. *Ecology* 68:1789–1809
- Smit C, Rietkerk M, Wassen MJ (2009) Inclusion of biotic stress (consumer pressure) alters predictions from the stress gradient hypothesis. *Journal of Ecology* 97:1215–1219
- Smith R, Tighe M, Reid N, Briggs S, Wilson B (2013) Effects of grazing, trenching and surface soil disturbance on ground cover in woody encroachment on the Cobar Pediplain, south-eastern Australia. *Journal of Arid Environments* 96:80–86
- Soliveres S, Van Der Plas F, Manning P, Prati D, Gossner MM, Renner SC, et al. (2016) Biodiversity at multiple trophic levels is needed for ecosystem multifunctionality. *Nature* 536:456–539
- Van Auker OW (2000) Shrub invasions of North American semiarid grasslands. *Annual Review of Ecology and Systematics* 31:197–215
- Watson VH, Ward CY (1970) Influence of intact tillers and height of cut on regrowth and carbohydrate reserves of dallisgrass (*Paspalum dilatatum* Poir.). *Crop Science* 10:474–476
- Wiedemann HT, Kelly PJ (2001) Turpentine (*Eremophila sturtii*) control by mechanical uprooting. *Rangeland Journal* 23:173–181

- Xiong X, Han X (2005) Spatial heterogeneity in soil carbon and nitrogen resources, caused by *Caragana microphylla*, in the thickening of semi-arid grassland, Inner Mongolia. *Acta Ecologica Sinica* 25:1678–1683
- Xiong X, Han X, Bai Y, Pan Q (2003) Increased distribution of *Caragana microphylla* in rangelands and its causes and consequences in Xilin River Basin. *Acta Prataculturae Sinica* 12:57–62
- Zavaleta ES, Hobbs RJ, Mooney HA (2001) Viewing invasive species removal in a whole-ecosystem context. *Trends in Ecology & Evolution* 16: 454–459
- Zhang Z, Wang SP, Nyren P, Jiang GM (2006) Morphological and reproductive response of *Caragana microphylla* to different stocking rates. *Journal of Arid Environments* 67:671–677
- Zhang H-Y, Lu X-T, Knapp AK, Hartmann H, Bai E, Wang X-B, Wang Z-W, Wang X-G, Yu Q, Han X-G (2018) Facilitation by leguminous shrubs increases along a precipitation gradient. *Functional Ecology* 32:203–213
- Zhao H, Zhou R, Su Y, Zhang H, Zhao L, Drake S (2007) Shrub facilitation of desert land restoration in the Horqin Sand Land of Inner Mongolia. *Ecological Engineering* 31:1–8

Coordinating Editor: Kari Veblen

Received: 17 April, 2019; First decision: 10 July, 2019; Revised: 10 July, 2019; Accepted: 22 May, 2019; First published online: 4 August, 2019